




OVER-THE-COUNTER MEDICINE SAFETY

UNDERSTANDING THE LABEL

The *Drug Facts* label helps you understand the medicines that you take and how to take them safely. All medicines should be taken with the direct supervision of a parent or trusted adult.



USES
Describes the symptoms that the medicine treats.



DIRECTIONS
Indicates the amount or dose of medicine to take, how often to take it, and how much you can take in one day.



INACTIVE INGREDIENTS
Ingredients not intended to treat your symptoms (e.g., preservatives, flavorings).



QUESTIONS OR COMMENTS?
Call the company if you have questions about a specific medicine.



ACTIVE INGREDIENTS
The ingredients in the medicine that make it work.



WARNINGS
Safety information, including side effects, the questions you should ask a doctor before taking the medicine, and which medicines to avoid using at the same time.



OTHER INFORMATION
How to store the medicine.

Drug Facts

Active ingredients

Medicine 100 mg	cough suppressant
Medicine 150 mg	nasal decongestant

Purpose

Uses Temporarily relieves:

- coughing due to minor throat and bronchial irritation
- nasal congestion

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product

- you may get drowsy
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and seek medical help right away if allergic reaction occurs.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- Tablet melts in mouth. Can be taken with or without water.

Age	Dose
adults and children 12 years and older	2 tablets every 12 hours; do not use more than 4 tablets in a 24-hour period
children 6 years to under 12 years	1 tablet every 12 hours; do not use more than 2 tablets in a 24-hour period
children under 6 years of age	ask a doctor

Other information

- store at 20°–25°C (68°–77°F)
- keep dry

Inactive ingredients

anhydrous citric acid, aspartame, magnesium stearate, maltodextrin, modified food starch, sodium bicarbonate, D&C yellow no.10


Questions or comments?

Call weekdays from 9 a.m. to 5 p.m. EST at 1-800-XXX-XXXX.

This is not an actual *Drug Facts* label.

MEASURE IT CORRECTLY

To get an exact measurement, it is important to only use the dosing device that is provided with the medicine. The *Drug Facts* label tells you how much medicine you should take based on your weight and/or age. Take only one medicine with the same kind of active ingredient at a time.



FACT: Household spoons come in various shapes and sizes, which means they hold different amounts of liquid.

ASK FOR HELP

Always talk with a parent or trusted adult to learn more about the responsible use of all medicines. Other resources include doctors, nurses, and pharmacists as well as your local poison control center.

In 2016, America's poison control centers managed about

630,000

medicine exposure cases involving children, tweens, and teens.¹

Poison Help
1-800-222-1222

DID YOU KNOW?

Poison control centers are staffed with experts, including doctors, nurses, and pharmacists who can help answer questions about medicines or can provide help over the phone if you come in contact with a poison. Calls are free and private.

¹Source: American Association of Poison Control Centers' National Poison Data System. Data covers ages 0–19.